

The Open Systems View and the Principle of Complete Positivity

(based on joint work with Stephan Hartmann)

Michael E. Cuffaro^{1,2}

¹Munich Center for Mathematical Philosophy, LMU Munich,
Michael.Cuffaro@lmu.de

²Generously supported by the Alexander von Humboldt Foundation and by the German Research Council (DFG)

September 23, 2023

European Philosophy of Science Association Biennial Meeting
Belgrade, Serbia

Bristol-Lancaster-Munich research group on open systems



Stephan Hartmann




James Ladyman



Sébastien Rivat



David Sloan



Karim Thébault



M. E. C.

M. E. C. and Hartmann, S. (eds.),
Open Systems: Physics, Metaphysics, and Methodology,
Oxford University Press (in preparation)

With chapters by:

- Emily Adlam, Luis C. Barbado and Časlav Brukner, Elena Castellani and Emilia Margoni, Eddy K. Chen, Elise Crull, M. E. C. and Stephan Hartmann, Richard Dawid, Gemma De las Cuevas, George Ellis, Doreen Fraser and Adam Koberinski, Sean Gryb and David Sloan, William L. Harper, Molly Kao, James Ladyman and Karim Thébault, Olimpia Lombardi, Wayne C. Myrvold, Daniele Oriti, Josh Quirke and Alistair Wilson, Katie Robertson, Karim Thébault, Lev Vaidman, and David Wallace.

Outline:

1. Standard quantum theory (**ST**) and the closed systems view
2. Reasons to worry about the closed systems view
3. The general quantum theory of open systems (**GT**) and the open systems view
4. The principle of complete positivity

Outline:

1. Standard quantum theory (**ST**) and the closed systems view
2. Reasons to worry about the closed systems view
3. The general quantum theory of open systems (**GT**) and the open systems view
4. The principle of complete positivity

The closed systems view

- (1) Metaphysical motivation: Isolated systems are the proper subject matter of science.
- (2) Associated with the methodology that models all phenomena fundamentally in terms of closed systems.

A “view” is similar to a “stance” (van Fraassen) but unlike a stance includes (1) in addition to (2).

Standard quantum theory (**ST**)

- Formulated in accordance with the closed systems view.
 - I.e., a given system's dynamics is generated by its Hamiltonian, which doesn't include any terms to reflect the system's interaction with something external.

Standard quantum theory (**ST**)

- Formulated in accordance with the closed systems view.
 - I.e., a given system's dynamics is generated by its Hamiltonian, which doesn't include any terms to reflect the system's interaction with something external.
- Physical state of a closed system, \mathcal{S} , is represented by a state vector, $|\psi\rangle$, in a Hilbert space for \mathcal{S} .
- Dynamics of $|\psi\rangle$ is unitary:

$$|\psi_t\rangle = U_t|\psi_0\rangle$$

Standard quantum theory (**ST**)

- Formulated in accordance with the closed systems view.
 - I.e., a given system's dynamics is generated by its Hamiltonian, which doesn't include any terms to reflect the system's interaction with something external.
- Physical state of a closed system, \mathcal{S} , is represented by a state vector, $|\psi\rangle$, in a Hilbert space for \mathcal{S} .
- Dynamics of $|\psi\rangle$ is unitary:

$$|\psi_t\rangle = U_t|\psi_0\rangle$$

Strictly speaking no system (except, perhaps, the whole universe) can really be isolated. How do we model open systems in **ST**?

Examples of quantum theories of open systems formulated in the framework of **ST**:

- Weisskopf-Wigner theory of spontaneous emissions
- Laser theory
- Quantum information theory

Examples of quantum theories of open systems formulated in the framework of **ST**:

- Weisskopf-Wigner theory of spontaneous emissions
- Laser theory
- Quantum information theory

Closed systems view of an open system:

- \mathcal{S} 's interaction with its environment is described in terms of its being coupled with a separate system \mathcal{E} such that $\mathcal{S} + \mathcal{E}$ form an isolated system.



Example: Deriving the Lindblad equation in **ST**

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.

Example: Deriving the Lindblad equation in **ST**

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.
 - Assume \mathcal{E} affects \mathcal{S} but not vice versa (weak coupling).

Example: Deriving the Lindblad equation in **ST**

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.
 - Assume \mathcal{E} affects \mathcal{S} but not vice versa (weak coupling).
 - Assume the future state of \mathcal{S} only depends on its present state (Markov approximation).

Example: Deriving the Lindblad equation in ST

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.
 - Assume \mathcal{E} affects \mathcal{S} but not vice versa (weak coupling).
 - Assume the future state of \mathcal{S} only depends on its present state (Markov approximation).
- Unitarily evolve the state, $|\Psi\rangle_{\mathcal{S}+\mathcal{E}}$, of the closed system $\mathcal{S} + \mathcal{E}$ and take the partial trace with respect to \mathcal{E} .

Example: Deriving the Lindblad equation in ST

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.
 - Assume \mathcal{E} affects \mathcal{S} but not vice versa (weak coupling).
 - Assume the future state of \mathcal{S} only depends on its present state (Markov approximation).
- Unitarily evolve the state, $|\Psi\rangle_{\mathcal{S}+\mathcal{E}}$, of the closed system $\mathcal{S} + \mathcal{E}$ and take the partial trace with respect to \mathcal{E} .
- Final state of \mathcal{S} : given by a reduced density operator:

$$\rho_{\mathcal{S}} = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

Example: Deriving the Lindblad equation in ST

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.
 - Assume \mathcal{E} affects \mathcal{S} but not vice versa (weak coupling).
 - Assume the future state of \mathcal{S} only depends on its present state (Markov approximation).
- Unitarily evolve the state, $|\Psi\rangle_{\mathcal{S}+\mathcal{E}}$, of the closed system $\mathcal{S} + \mathcal{E}$ and take the partial trace with respect to \mathcal{E} .
- Final state of \mathcal{S} : given by a reduced density operator:

$$\rho_{\mathcal{S}} = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

- Yields (effective) non-unitary dynamics for \mathcal{S} :

$$\dot{\rho}_{\mathcal{S}} = (\mathcal{L}_u + \mathcal{L}_{n-u}) \rho_{\mathcal{S}},$$

$$\text{with: } \mathcal{L}_{n-u} \rho = \frac{1}{2} \sum_i \left([L_i \rho, L_i^\dagger] + [L_i, \rho L_i^\dagger] \right),$$

where the L_i are (bounded) operators.

Outline:

1. Standard quantum theory (**ST**) and the closed systems view
2. Reasons to worry about the closed systems view
3. The general quantum theory of open systems (**GT**) and the open systems view
4. The principle of complete positivity

Prima facie reasons to worry about the closed systems view:

- Standard models of cosmology describe the universe as closed but are often based on strong idealizations introduced only to simplify the mathematics (Smeenk & Ellis, 2017, Sec. 1.1).

Prima facie reasons to worry about the closed systems view:

- Standard models of cosmology describe the universe as closed but are often based on strong idealizations introduced only to simplify the mathematics (Smeenk & Ellis, 2017, Sec. 1.1).
 - Thus, although our best cosmological models describe our universe as a closed system, this does not necessarily mean that our universe actually is a closed system (see also Gryb & Sloan, 2021; Sloan, 2018).

Prima facie reasons to worry about the closed systems view:

- Standard models of cosmology describe the universe as closed but are often based on strong idealizations introduced only to simplify the mathematics (Smeenk & Ellis, 2017, Sec. 1.1).
 - Thus, although our best cosmological models describe our universe as a closed system, this does not necessarily mean that our universe actually is a closed system (see also Gryb & Sloan, 2021; Sloan, 2018).
- Black hole physics gives us (prima facie) reasons to motivate the idea that the dynamics of certain systems are fundamentally open (Hawking, 1976).

Prima facie reasons to worry about the closed systems view:

- Standard models of cosmology describe the universe as closed but are often based on strong idealizations introduced only to simplify the mathematics (Smeenk & Ellis, 2017, Sec. 1.1).
 - Thus, although our best cosmological models describe our universe as a closed system, this does not necessarily mean that our universe actually is a closed system (see also Gryb & Sloan, 2021; Sloan, 2018).
- Black hole physics gives us (prima facie) reasons to motivate the idea that the dynamics of certain systems are fundamentally open (Hawking, 1976).
- Global unitary evolution is hard to square with important recent approaches to quantum gravity (Oriti, 2021, sec. 3.1).

A more principled reason to worry about the closed systems view:

- In practice we strictly speaking only ever apply quantum theory to open systems (except in cosmology).

A more principled reason to worry about the closed systems view:

- In practice we strictly speaking only ever apply quantum theory to open systems (except in cosmology).
- Although this typically involves the methodological assumptions associated with the closed systems view,

A more principled reason to worry about the closed systems view:

- In practice we strictly speaking only ever apply quantum theory to open systems (except in cosmology).
- Although this typically involves the methodological assumptions associated with the closed systems view, **it isn't the dynamics of $\mathcal{S} + \mathcal{E}$, but the dynamics of \mathcal{S} , that we take ourselves to have successfully described when we do this.**

A more principled reason to worry about the closed systems view:

- In practice we strictly speaking only ever apply quantum theory to open systems (except in cosmology).
- Although this typically involves the methodological assumptions associated with the closed systems view, **it isn't the dynamics of $\mathcal{S} + \mathcal{E}$, but the dynamics of \mathcal{S} , that we take ourselves to have successfully described when we do this.**
- Thus there is a clear **empirical motivation to extrapolate** from the dynamics of open systems rather than from the dynamics of closed systems.

Cf. Newton's 4th rule of reasoning: "In experimental philosophy, propositions gathered from phenomena by induction should be considered either exactly or very nearly true notwithstanding any contrary hypotheses, until yet other phenomena make such propositions either more exact or liable to exceptions." (Quoted in Harper 2011, ch. 7)

Even deeper reasons to worry about the closed systems view:

Approaches to the interpretation of **ST** that take it to be complete
(in some sense):

Even deeper reasons to worry about the closed systems view:

Approaches to the interpretation of **ST** that take it to be complete (in some sense):

- “(neo-)Everettian” approaches: **ST** provides us with a complete description of physical reality.

Even deeper reasons to worry about the closed systems view:

Approaches to the interpretation of **ST** that take it to be complete (in some sense):

- “(neo-)Everettian” approaches: **ST** provides us with a complete description of physical reality.
- “(neo-)Bohrian” approaches: **ST** provides us with all of the resources one needs to describe any given probabilistic physical phenomenon to whatever level of detail one would like.

Even deeper reasons to worry about the closed systems view:

Approaches to the interpretation of **ST** that take it to be complete (in some sense):

- “(neo-)Everettian” approaches: **ST** provides us with a complete description of physical reality.
- “(neo-)Bohrian” approaches: **ST** provides us with all of the resources one needs to describe any given probabilistic physical phenomenon to whatever level of detail one would like.

Both (neo-)Everettian and (neo-)Bohrian approaches are ontologically committed to open systems (or so we argue).

$$\begin{aligned} |\psi\rangle_{\mathcal{S}} &= \alpha|b_1^+\rangle + \beta|b_1^-\rangle \\ &= \alpha'|b_2^+\rangle + \beta'|b_2^-\rangle. \end{aligned}$$

What does this mean on a (neo-)Bohrian interpretation?

- **Coupling the degrees of freedom of \mathcal{S} to those of a further system \mathcal{M}** will yield a collection of unitarily-related conditional probability distributions over the possible outcomes of an assessment of \mathcal{M} as described with respect to a particular basis b_m .

“In the treatment of atomic problems, actual calculations are most conveniently carried out with the help of a Schrödinger state function, from which the statistical laws governing observations obtainable under specified conditions can be deduced by definite mathematical operations. It must be recognized, however, that we are here dealing with a purely symbolic procedure, **the unambiguous physical interpretation of which in the last resort requires a reference to a complete experimental arrangement.**” (Bohr, 1958, pp. 392–393, our emphasis).

(neo-)Bohrian interpretation:

- Notice that \mathcal{S} is conceived of here as an open system (even when its state is described by a state vector), **but since open systems dynamics are not fundamental in \mathbf{ST} , we require a larger Hilbert space** (including the degrees of freedom of both \mathcal{S} and \mathcal{M}) to represent it as such.

(neo-)Bohrian interpretation:

- Notice that \mathcal{S} is conceived of here as an open system (even when its state is described by a state vector), **but since open systems dynamics are not fundamental in ST**, we require a **larger Hilbert space** (including the degrees of freedom of both \mathcal{S} and \mathcal{M}) to represent it as such.
- **ST is about open systems**, on a (neo-)Bohrian interpretation, despite being formulated from the closed systems view (which it inherits from classical mechanics).

The Everett interpretation:

- **ST** is as much about subsystems described by density operators as it is about composite systems described by state vectors (see, e.g., Wallace and Timpson's (2010): "Spacetime state realism.")

The Everett interpretation:

- **ST** is as much about subsystems described by density operators as it is about composite systems described by state vectors (see, e.g., Wallace and Timpson's (2010): "Spacetime state realism.")
- More principled argument: Corresponding to any given probability distribution over pure states of \mathcal{B} (i.e., to any given density operator for \mathcal{B}), one can always find a pure state of some larger system $\mathcal{A} + \mathcal{B}$ from which that probability distribution can be derived.
- All such purifications are "essentially the same" (D'Ariano et al., 2017, p. 171).

The Everett interpretation:

- **ST** is as much about subsystems described by density operators as it is about composite systems described by state vectors (see, e.g., Wallace and Timpson's (2010): "Spacetime state realism.")
- More principled argument: Corresponding to any given probability distribution over pure states of \mathcal{B} (i.e., to any given density operator for \mathcal{B}), one can always find a pure state of some larger system $\mathcal{A} + \mathcal{B}$ from which that probability distribution can be derived.
- All such purifications are "essentially the same" (D'Ariano et al., 2017, p. 171).
- $\rho_{\mathcal{S}}$, expressed as a decoherent mixture of various states corresponding to the elements of an eigenbasis of $\rho_{\mathcal{M}}$ is just as objective a description of everything there is, relative to the degrees of freedom included in our representation of \mathcal{S} and to the given eigenbasis, as the universal state vector $|\Psi\rangle_{\mathcal{S}+\mathcal{M}}$.

The Everett interpretation:

Ultimately, what is the Everett interpretation committed to?

The Everett interpretation:

Ultimately, what is the Everett interpretation committed to?

- Fundamental unitarity?

The Everett interpretation:

Ultimately, what is the Everett interpretation committed to?

- Fundamental unitarity?
 - Not necessarily!
 - Taking the non-unitary dynamics of a density operator to be fundamental, **even in the case where it represents the universe as a whole**, is consistent with the Everett interpretation (see, e.g., Wallace 2012, sec. 10.5).

The Everett interpretation:

Ultimately, what is the Everett interpretation committed to?

- Fundamental unitarity?
 - Not necessarily!
 - Taking the non-unitary dynamics of a density operator to be fundamental, **even in the case where it represents the universe as a whole**, is consistent with the Everett interpretation (see, e.g., Wallace 2012, sec. 10.5).
 - Ultimately the Everett interpretation is committed to **quantum theory**, not necessarily the closed systems view of quantum theory.

Outline:

1. Standard quantum theory (**ST**) and the closed systems view
2. Reasons to worry about the closed systems view
3. The general quantum theory of open systems (**GT**) and the open systems view
4. The principle of complete positivity

The open systems view

- (1) Metaphysical motivation: The proper subject matter of science is systems that are, in general, open.
- (2) Associated with the methodology according to which one need not model phenomena in terms of closed systems. The influence of the environment on a system is represented fundamentally in terms the dynamical equations that we take to govern its evolution.



General Quantum Theory of Open Systems (GT)

- Formulated in accordance with the open systems view.
 - The environment is not represented as a separate system; its influence is represented in the dynamical equations that govern the evolution of \mathcal{S} .

General Quantum Theory of Open Systems (GT)

- Formulated in accordance with the open systems view.
 - The environment is not represented as a separate system; its influence is represented in the dynamical equations that govern the evolution of \mathcal{S} .
- Physical state of \mathcal{S} represented by a density operator, ρ
- Time-evolution of ρ is governed by a dynamical map, Λ_t :

$$\Lambda_t \rho_0 = \rho_t$$

- Λ_t acts on the state space of \mathcal{S} (not on $\mathcal{S} + \mathcal{E}$).

General derivation of the Lindblad equation in **GT**

- The family of dynamical maps for a given system form a **semigroup** (in particular, no inverse).

General derivation of the Lindblad equation in **GT**

- The family of dynamical maps for a given system form a **semigroup** (in particular, no inverse).
- The dynamics of \mathcal{S} is assumed to be **Markovian**: probabilities at t_1 are uniquely determined given probabilities at t_0 , but in general not vice versa.

General derivation of the Lindblad equation in **GT**

- The family of dynamical maps for a given system form a **semigroup** (in particular, no inverse).
- The dynamics of \mathcal{S} is assumed to be **Markovian**: probabilities at t_1 are uniquely determined given probabilities at t_0 , but in general not vice versa.
- The dynamics of \mathcal{S} is assumed to be **completely positive**.

General derivation of the Lindblad equation in GT

- The family of dynamical maps for a given system form a **semigroup** (in particular, no inverse).
- The dynamics of \mathcal{S} is assumed to be **Markovian**: probabilities at t_1 are uniquely determined given probabilities at t_0 , but in general not vice versa.
- The dynamics of \mathcal{S} is assumed to be **completely positive**.
- \mathcal{S} is assumed to evolve **continuously** through time.
 - As $t \rightarrow 0$, $\Lambda_t \rho \approx \rho$

General derivation of the Lindblad equation in GT

- The family of dynamical maps for a given system form a **semigroup** (in particular, no inverse).
- The dynamics of \mathcal{S} is assumed to be **Markovian**: probabilities at t_1 are uniquely determined given probabilities at t_0 , but in general not vice versa.
- The dynamics of \mathcal{S} is assumed to be **completely positive**.
- \mathcal{S} is assumed to evolve **continuously** through time.
 - As $t \rightarrow 0$, $\Lambda_t \rho \approx \rho$

Consequence:

$$\dot{\rho}_{\mathcal{S}} = (\mathcal{L}_u + \mathcal{L}_{n-u}) \rho_{\mathcal{S}}$$

- Non-unitary dynamics in general; unitary dynamics as a special case.

There is more to **GT** than the Lindblad equation. We can consider relaxing some of its assumptions. For instance,

- Markov (Barandes, 2023)
- Semigroup, continuity (Wolf & Cirac, 2008).
- Complete positivity: The focus of the rest of this talk.

Outline:

1. Standard quantum theory (**ST**) and the closed systems view
2. Reasons to worry about the closed systems view
3. The general quantum theory of open systems (**GT**) and the open systems view
4. The principle of complete positivity

Complete positivity:

Our claim:

- Complete positivity is an expression of the closed systems view. “Not completely positive” (NCP) maps do not make physical sense on the closed systems view.
- But on the open systems view, one should deny complete positivity the status of a fundamental physical principle.

ρ_S (“density operator”):

- Probabilistic generalisation of a state vector:

$$\rho_S = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

ρ_S (“density operator”):

- Probabilistic generalisation of a state vector:

$$\rho_S = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

- Positive semidefinite operator of unit trace
 - non-negative probabilities

ρ_S (“density operator”):

- Probabilistic generalisation of a state vector:

$$\rho_S = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

- Positive semidefinite operator of unit trace
 - non-negative probabilities
- Dynamics:
 - Isolated system: $\rho_S \mapsto \mathbf{U}\rho_S\mathbf{U}^\dagger$

ρ_S (“density operator”):

- Probabilistic generalisation of a state vector:

$$\rho_S = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

- Positive semidefinite operator of unit trace
 - non-negative probabilities
- Dynamics:
 - Isolated system: $\rho_S \mapsto \mathcal{U}\rho_S\mathcal{U}^\dagger$
 - Compound system (product state):
 $\rho_{S+\mathcal{E}} \mapsto \mathcal{U}\rho_{S+\mathcal{E}}\mathcal{U}^\dagger = \mathcal{U}\rho_S \otimes \rho_{\mathcal{E}}\mathcal{U}^\dagger$

$\rho_{\mathcal{S}}$ (“density operator”):

- Probabilistic generalisation of a state vector:

$$\rho_{\mathcal{S}} = \sum_i p_i |\psi_i\rangle \langle \psi_i|$$

- Positive semidefinite operator of unit trace
 - non-negative probabilities
- Dynamics:
 - Isolated system: $\rho_{\mathcal{S}} \mapsto \mathbf{U}\rho_{\mathcal{S}}\mathbf{U}^\dagger$
 - Compound system (product state):
 $\rho_{\mathcal{S}+\mathcal{E}} \mapsto \mathbf{U}\rho_{\mathcal{S}+\mathcal{E}}\mathbf{U}^\dagger = \mathbf{U}\rho_{\mathcal{S}} \otimes \rho_{\mathcal{E}}\mathbf{U}^\dagger$
- State change of \mathcal{S} in the presence of \mathcal{E} :
 $\rho_{\mathcal{S}} \mapsto \text{tr}_{\mathcal{E}}(\mathbf{U}\rho_{\mathcal{S}} \otimes \rho_{\mathcal{E}}\mathbf{U}^\dagger) = \rho'_{\mathcal{S}} = \Lambda\rho_{\mathcal{S}}$

$$\rho_S \mapsto \text{tr}_E(\mathbf{U}\rho_S \otimes \rho_E \mathbf{U}^\dagger) = \Lambda\rho_S$$


$$\rho_S \mapsto \text{tr}_E(\mathbf{U}\rho_S \otimes \rho_E \mathbf{U}^\dagger) = \Lambda\rho_S$$

- Λ : maps ρ_S to ρ'_S , $\Rightarrow \Lambda$ must be a positive map.

$$\rho_{\mathcal{S}} \mapsto \text{tr}_{\mathcal{E}}(\mathbf{U}\rho_{\mathcal{S}} \otimes \rho_{\mathcal{E}}\mathbf{U}^{\dagger}) = \Lambda\rho_{\mathcal{S}}$$

- Λ : maps $\rho_{\mathcal{S}}$ to $\rho'_{\mathcal{S}}$, $\Rightarrow \Lambda$ must be a positive map.

Complete positivity:




Consider the system $\mathcal{S} + \mathcal{W}_n$ evolving in the presence of \mathcal{E} , such that:

- \mathcal{W}_n : A system of dimensionality n not currently interacting with \mathcal{S} , but which may have interacted with it in the past.
- \mathcal{W}_n evolves trivially.

$$\rho_S \mapsto \text{tr}_E(\mathbf{U}\rho_S \otimes \rho_E \mathbf{U}^\dagger) = \Lambda\rho_S$$

- Λ : maps ρ_S to ρ'_S , $\Rightarrow \Lambda$ must be a positive map.

Complete positivity:



Consider the system $\mathcal{S} + \mathcal{W}_n$ evolving in the presence of \mathcal{E} , such that:

- \mathcal{W}_n : A system of dimensionality n not currently interacting with \mathcal{S} , but which may have interacted with it in the past.
- \mathcal{W}_n evolves trivially.


Problem: Requiring that Λ be positive on \mathcal{S} does not guarantee that $\Lambda \otimes I_n$ is positive on $\mathcal{S} + \mathcal{W}_n$.

- I.e., $\Lambda \otimes I_n$ will map some of the states in $\mathcal{H}_{\mathcal{S} + \mathcal{W}_n}$ to unphysical states (i.e., that yield negative probabilities for the results of measurements on $\mathcal{S} + \mathcal{W}_n$).

$$\rho_S \mapsto \text{tr}_E(\mathbf{U}\rho_S \otimes \rho_E \mathbf{U}^\dagger) = \Lambda\rho_S$$

- Λ : maps ρ_S to ρ'_S , $\Rightarrow \Lambda$ must be a positive map.

Complete positivity:



Consider the system $S + \mathcal{W}_n$ evolving in the presence of E , such that:

- \mathcal{W}_n : A system of dimensionality n not currently interacting with S , but which may have interacted with it in the past.
- \mathcal{W}_n evolves trivially.

Problem: Requiring that Λ be positive on S does not guarantee that $\Lambda \otimes I_n$ is positive on $S + \mathcal{W}_n$.

- I.e., $\Lambda \otimes I_n$ will map some of the states in $\mathcal{H}_{S+\mathcal{W}_n}$ to unphysical states (i.e., that yield negative probabilities for the results of measurements on $S + \mathcal{W}_n$).

Solution: Require that $\Lambda \otimes I_n$ be positive for all \mathcal{W}_n .

“One may reasonably doubt this argument. It is very powerful magic: \mathcal{W} sits apart from $\mathcal{S} + \mathcal{E}$ and does absolutely nothing; by doing so, it forces the motion of \mathcal{S} to be completely positive with dramatic physical consequences ...” (Pechukas, 1994).

More concrete reasons to be skeptical (Shaji & Sudarshan, 2005):

Suppose, on the one hand, that \mathcal{S} and \mathcal{W}_n are not entangled:

- A positive but not completely positive map on \mathcal{S} 's state space will always entail a valid evolution for $\mathcal{S} + \mathcal{W}_n$ under the remaining assumptions.

More concrete reasons to be skeptical (Shaji & Sudarshan, 2005):

Suppose, on the one hand, that \mathcal{S} and \mathcal{W}_n are not entangled:

- A positive but not completely positive map on \mathcal{S} 's state space will always entail a valid evolution for $\mathcal{S} + \mathcal{W}_n$ under the remaining assumptions.

Suppose, on the other hand, that \mathcal{S} and \mathcal{W}_n are entangled:

- \mathcal{W}_n should really be thought of as part of the environment, \mathcal{E} .

More concrete reasons to be skeptical (Shaji & Sudarshan, 2005):

Suppose, on the one hand, that \mathcal{S} and \mathcal{W}_n are not entangled:

- A positive but not completely positive map on \mathcal{S} 's state space will always entail a valid evolution for $\mathcal{S} + \mathcal{W}_n$ under the remaining assumptions.

Suppose, on the other hand, that \mathcal{S} and \mathcal{W}_n are entangled:

- \mathcal{W}_n should really be thought of as part of the environment, \mathcal{E} .
- Important because it can be shown (Jordan et al., 2004, pp. 13–14) that the contracted dynamics of a system \mathcal{S} in the presence of \mathcal{E} is describable by a completely positive map only if \mathcal{S} is initially not entangled with \mathcal{E} .

More concrete reasons to be skeptical (Shaji & Sudarshan, 2005):

Suppose, on the one hand, that \mathcal{S} and \mathcal{W}_n are not entangled:

- A positive but not completely positive map on \mathcal{S} 's state space will always entail a valid evolution for $\mathcal{S} + \mathcal{W}_n$ under the remaining assumptions.

Suppose, on the other hand, that \mathcal{S} and \mathcal{W}_n are entangled:

- \mathcal{W}_n should really be thought of as part of the environment, \mathcal{E} .
- Important because it can be shown (Jordan et al., 2004, pp. 13–14) that the contracted dynamics of a system \mathcal{S} in the presence of \mathcal{E} is describable by a completely positive map only if \mathcal{S} is initially not entangled with \mathcal{E} .
- If we enforce complete positivity then it would seem to follow that no valid physical description of the dynamics of \mathcal{S} can be given when it is initially entangled with \mathcal{E} .

Not completely positive (NCP) maps:

It is precisely for setups like these that an NCP map will make sense.

- When \mathcal{S} and \mathcal{E} are entangled, then it follows that it is impossible for \mathcal{S} to be (for instance) in a pure state, or in general in any state that is not a valid partial trace over the combined state of $\mathcal{S} + \mathcal{E}$.

Not completely positive (NCP) maps:

It is precisely for setups like these that an NCP map will make sense.

- When \mathcal{S} and \mathcal{E} are entangled, then it follows that it is impossible for \mathcal{S} to be (for instance) in a pure state, or in general in any state that is not a valid partial trace over the combined state of $\mathcal{S} + \mathcal{E}$.
- Such states that are outside of the 'compatibility domain' of an NCP map will be ill-described by it.

Not completely positive (NCP) maps:

It is precisely for setups like these that an NCP map will make sense.

- When \mathcal{S} and \mathcal{E} are entangled, then it follows that it is impossible for \mathcal{S} to be (for instance) in a pure state, or in general in any state that is not a valid partial trace over the combined state of $\mathcal{S} + \mathcal{E}$.
- Such states that are outside of the 'compatibility domain' of an NCP map will be ill-described by it.
- But as long as such a map is completely positive **in relation to all of the actually possible states of \mathcal{S}** in a given setup, it seems that there is no reason not to use it to describe the dynamics of \mathcal{S} (Cuffaro & Myrvold, 2013, sec. 5).

A better argument for imposing complete positivity as a fundamental physical principle:

*A system-theoretic description of an open system has to be considered as phenomenological; **the requirement that it should be derivable from the fundamental automorphic dynamics of a closed system** implies that the dynamical map of an open system has to be completely positive. (Raggio & Primas, 1982, p. 435, our emphasis).*

Deriving the Lindblad equation in $\mathcal{S}\mathcal{T}$

- We consider a particular open system \mathcal{S} such that the following assumptions can be taken to hold.
 - Assume \mathcal{E} affects \mathcal{S} but not vice versa (weak coupling).
 - Assume the future state of \mathcal{S} only depends on its present state (Markov approximation).

- Unitarily evolve the state, $|\Psi\rangle_{\mathcal{S}+\mathcal{E}}$, of the closed system $\mathcal{S} + \mathcal{E}$ and take the partial trace with respect to \mathcal{E} .

- Final state of \mathcal{S} : given by a reduced density operator:

$$\rho_{\mathcal{S}} = \sum_i p_i |\psi_i\rangle\langle\psi_i|$$

- Yields (effective) non-unitary dynamics for \mathcal{S} :

$$\dot{\rho}_{\mathcal{S}} = (\mathcal{L}_u + \mathcal{L}_{n-u}) \rho_{\mathcal{S}},$$

$$\text{with: } \mathcal{L}_{n-u} \rho = \frac{1}{2} \sum_i \left([L_i \rho, L_i^\dagger] + [L_i, \rho L_i^\dagger] \right),$$

where the L_i are (bounded) operators.

- Guaranteed by Stinespring's dilation theorem (Stinespring, 1955), which assumes complete positivity.

Since, in **GT**, open systems dynamics are taken to be fundamental,

Since, in **GT**, open systems dynamics are taken to be fundamental,

- there's no need to understand this in terms of a contraction of the dynamics of a larger closed system.

Since, in **GT**, open systems dynamics are taken to be fundamental,

- there's no need to understand this in terms of a contraction of the dynamics of a larger closed system.
- It follows that the dynamical maps governing an open system do not, in general, need to be completely positive.

Since, in **GT**, open systems dynamics are taken to be fundamental,

- there's no need to understand this in terms of a contraction of the dynamics of a larger closed system.
- It follows that the dynamical maps governing an open system do not, in general, need to be completely positive.
- This makes it possible to describe the evolution of even the universe as a whole (assuming one takes that to make sense given one's interpretation of QM) as if it were initially a subsystem of an entangled system.

Since, in **GT**, **open systems dynamics are taken to be fundamental**,

- there's no need to understand this in terms of a contraction of the dynamics of a larger closed system.
- It follows that the dynamical maps governing an open system do not, in general, need to be completely positive.
- This makes it possible to describe the evolution of even the universe as a whole (assuming one takes that to make sense given one's interpretation of QM) as if it were initially a subsystem of an entangled system.

So **GT**, unlike ST, allows for **fundamental non-unitary evolution**, and is, in this sense, a more general dynamical framework than **ST** (despite not adding anything to the Hilbert space formalism).

~~A system-theoretic description of an open system has to be considered as phenomenological; the requirement that it should be derivable from the fundamental automorphic dynamics of a closed system implies that the dynamical map of an open system has to be completely positive. (Raggio & Primas, 1982, p. 435).~~

Conclusion: We should reject complete positivity as a fundamental physical principle.

For more on the open systems view, see [arXiv:2112.11095](#), [arXiv:2305.00378](#), and our forthcoming book:

M. E. C. and Hartmann, S. (eds.),
Open Systems: Physics, Metaphysics, and Methodology,
Oxford University Press (in preparation)

With chapters by:

- Emily Adlam, Luis C. Barbado and Časlav Brukner, Elena Castellani and Emilia Margoni, Eddy K. Chen, Elise Crull, M. E. C. and Stephan Hartmann, Richard Dawid, Gemma De las Cuevas, George Ellis, Doreen Fraser and Adam Koberinski, Sean Gryb and David Sloan, William L. Harper, Molly Kao, James Ladyman and Karim Thébault, Olimpia Lombardi, Wayne C. Myrvold, Daniele Oriti, Josh Quirke and Alistair Wilson, Katie Robertson, Karim Thébault, Lev Vaidman, and David Wallace.

Extra slides

What about many worlds?

$$\rho = p |\psi_1\rangle\langle\psi_1| + (1 - p) |\psi_2\rangle\langle\psi_2|.$$

- The fact that the different terms, $|\psi_1\rangle\langle\psi_1|$ and $|\psi_2\rangle\langle\psi_2|$ are by definition **decoherent** makes it unproblematic, irrespective of whether ρ evolves unitarily, to identify them with independently evolving worlds.
- This is even clearer than the FAPP story one needs to give in the pure state, unitary, case.

A post-quantum theory of classical gravity?

Jonathan Oppenheim

We present a consistent theory of classical gravity coupled to quantum field theory. The dynamics is linear in the density matrix, completely positive and trace-preserving, and reduces to Einstein's theory of general relativity in the classical limit. As a result, the dynamics doesn't suffer from the pathologies of the semi-classical theory based on expectation values. The assumption that general relativity is classical necessarily modifies the dynamical laws of quantum mechanics -- the theory must be fundamentally stochastic in both the metric degrees of freedom and in the quantum matter fields. This allows it to evade several no-go theorems purporting to forbid classical-quantum interactions. The measurement postulate of quantum mechanics is not needed since the interaction of the quantum degrees of freedom with classical space-time necessarily causes decoherence. We first derive the most general form of classical-quantum dynamics and consider realisations which have as its limit deterministic classical Hamiltonian evolution. The formalism is then applied to quantum field theory interacting with the classical space-time metric. One can view the theory as fundamental or as an effective theory useful for computing the back-reaction of quantum fields on geometry.

Comments: "It's very difficult to find a black cat in a dark room, especially if there is no cat."

Subjects: **High Energy Physics - Theory (hep-th)**; Quantum Physics (quant-ph)

Cite as: arXiv:1811.03116 [hep-th]

(or arXiv:1811.03116v2 [hep-th] for this version)

<https://doi.org/10.48550/arXiv.1811.03116> 

An early attempt to couple classical gravity to quantum mechanics was made in [51] using the Aleksandrov-Gerasimenko bracket [52, 53]

$$\frac{\partial \sigma}{\partial t} \stackrel{?}{=} -i[\mathbf{H}, \sigma] + \frac{1}{2}(\{\mathbf{H}, \sigma\} - \{\sigma, \mathbf{H}\}) \quad (4)$$

However, the dynamics it generates leads to negative probabilities [51, 54]. Unless we are prepared to modify the Born rule, or equivalently, our interpretation of the density matrix, any dynamics must be trace-preserving (TP) to conserve probability, and be *completely positive* (CP), for probabilities to remain positive. So, although the dynamics of Equation (4) makes a frequent appearance in attempts to couple quantum and classical degrees of freedom [55-57] (c.f. [58]), and while it may give insight in some regimes, it cannot serve as a fundamental theory. Other attempts, such as using the Schrödinger-Newton equation [59, 60] (c.f. [61, 62]), suffer from the problem that the equations are non-linear in the density matrix, and thus will lead to superluminal signalling [63] and a breakdown of the statistical mechanical interpretation of the density matrix [7].

Works Cited I

- Barandes, J. A. (2023). The stochastic-quantum correspondence. arXiv:2302.10778.
- Bohr, N. (1958). Quantum physics and philosophy. In R. Klíbanký (Ed.) *Philosophy in the Mid-Century: A Survey*, (pp. 308–314). Firenze: La Nuova Italia Editrice. Page reference to reprint in: J. Kalckar, Ed., *Niels Bohr: Collected Works*, Vol. 7 (pp. 385–394). Amsterdam: Elsevier.
- Cuffaro, M. E., & Myrvold, W. C. (2013). On the debate concerning the proper characterisation of quantum dynamical evolution. *Philosophy of Science*, 80, 1125–1136.
- D'Ariano, G. M., Chiribella, G., & Perinotti, P. (2017). *Quantum Theory from First Principles: An Informational Approach*. Cambridge: Cambridge University Press. 2019 paperback edition.
- Gryb, S., & Sloan, D. (2021). When scale is surplus. *Synthese*, 199, 14769–14820.
- Harper, W. (2011). *Isaac Newton's Scientific Method: Turning Data Into Evidence*. Oxford: Oxford University Press.
- Hawking, S. W. (1976). Breakdown of predictability in gravitational collapse. *Physical Review D*, 14, 2460–2473.
- Jordan, T. F., Shaji, A., & Sudarshan, E. C. G. (2004). Dynamics of initially entangled open quantum systems. *Physical Review A*, 70, 052110.
- Oriti, D. (2021). The complex timeless emergence of time in quantum gravity. In P. Harris, & R. Lestienne (Eds.) *Time and Science*. World Scientific. Forthcoming.

Works Cited II

- Pechukas, P. (1994). Reduced dynamics need not be completely positive. *Physical Review Letters*, *73*, 1060–1062.
- Raggio, G. A., & Primas, H. (1982). Remarks on “On completely positive maps in generalized quantum dynamics”. *Foundations of Physics*, *12*, 433–435.
- Shaji, A., & Sudarshan, E. C. G. (2005). Who's afraid of not completely positive maps? *Physics Letters A*, *341*, 48–54.
- Sloan, D. (2018). Dynamical similarity. *Physical Review D*, *97*, 123541.
- Smeenk, C., & Ellis, G. (2017). Philosophy of cosmology. In E. N. Zalta (Ed.) *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, Winter 2017 ed.
- Stinespring, W. F. (1955). Positive functions on C^* -algebras. *Proceedings of the American Mathematical Society*, *6*, 211.
- Wallace, D. (2012). *The Emergent Multiverse*. Oxford: Oxford University Press.
- Wallace, D., & Timpson, C. G. (2010). Quantum mechanics on spacetime I: Spacetime state realism. *The British Journal for the Philosophy of Science*, *61*, 697–727.
- Wolf, M. M., & Cirac, I. (2008). Dividing quantum channels. *Communications in Mathematical Physics*, *279*, 147–168.